

Product Information

Bovine Aorta Endothelial Cells

Catalog Number	10BO-001	Cell Number	0.5 million cells/vial
Species	<i>Bos taurus</i>	Storage Temperature	Liquid Nitrogen

Description

Bovine Aorta Endothelial Cells (BAECs) line the vessel wall of aorta, and are constantly exposed to high hemodynamic forces. They produce endothelium-derived substances regulating vasoconstriction and vessel growth ^[1]. BAECs also modulate the expression of cellular adhesion molecules to control and fine-tune inflammatory responses and fibrinolysis ^[2]. These physiological properties allow BAEC cultures to be widely used in the study of mechanisms for endothelium dysfunction, pathogenesis of vascular diseases and atherosclerosis, and the development of novel disease treatments.

iXCells Biotechnologies offers BAECs from bovine aorta. These cells are negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast, and fungi. BAECs are guaranteed to maintain at least 16 passages.




Figure 1. Bovine Aorta Endothelial Cells (phase contrast).

Product Details

Tissue	Bovine aorta
Package Size	0.5 million cells/vial
Passage Number	P2
Shipped	Cryopreserved
Storage	Liquid nitrogen
Growth Properties	Adherent
Media	Endothelial Cell Growth Medium (Cat# MD-0010)

Protocols

Thawing of Frozen Cells

1. Upon receipt of the frozen cells, it is recommended to thaw the cells and initiate the culture immediately in order to retain the highest cell viability.
2. To thaw the cells, put the vial in 37°C water bath with gentle agitation for ~1 minute. Keep the cap out of water to minimize the risk of contamination.
3. Pipette the cells into a 15ml conical tube with 5ml fresh Endothelial Cell Growth Medium (Cat# MD-0010).
4. Centrifuge at 1000rpm (~220g) for 5 minutes under room temperature.
5. Remove the supernatant and resuspend the cells in Endothelial Cell Growth Medium.
6. Culture the cell in T75 flask.

Safety Precaution: *it is highly recommended that protective gloves and clothing should be used when handling frozen vials.*

Standard Culture Procedure

1. BAECs can be cultured in Endothelial Cell Growth Medium (Cat# MD-0010).
2. When cells reach ~80-90% confluence, remove the medium, and wash once with sterile PBS (5ml/T75 flask).
3. Add ~2.5ml of 0.25% Trypsin-EDTA to the flask and incubate for ~3 minutes at 37°C. Neutralize the enzyme by adding 2-3 volumes of cell culture medium.
4. Centrifuge 1,000 rpm (~220g) for 5min and resuspend the cells in desired volume of medium.
5. Seed new culture vessels at 5×10^3 cells/cm².

References

- [1] Ando J, and Kamiya A. Flow-dependent regulation of gene expression in vascular endothelial cells. Heart J. 1996; 37:19-32.
- [2] Liu JW, Wei DZ, etc. Enhancement of fibrinolytic activity of bovine aortic endothelial cells by ginsenoside Rb2. Acta Pharmacol sin 2003; 24: 102-108.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While iXCells Biotechnologies uses reasonable efforts to include accurate and up-to-date information on this product sheet, we make no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. iXCells Biotechnologies does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. iXCells Biotechnologies is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, iXCells Biotechnologies is not liable for damages arising from the misidentification or misrepresentation of cultures.

© iXCells Biotechnologies 2015. All rights reserved.